



血糖値 (HbA1c) を記録しましょう

※HbA1c値はNGSP値で表記

検査日											
検査機関											
糖尿病診療ガイドライン2013	 合併症の危険がさらに大きくなる！ 	HbA1c (JDS値)									
		9.4 (9.0)									
		9.2 (8.8)									
		9.0 (8.6)									
		8.8 (8.4)									
		8.6 (8.2)									
		8.4 (8.0)									
		8.3 (7.9)									
		8.2 (7.8)									
		8.1 (7.7)									
		8.0% (7.6)									
		7.9 (7.5)									
	7.8 (7.4)										
	7.7 (7.3)										
	7.6 (7.2)										
	7.5 (7.1)										
	7.4 (7.0)										
	7.3 (6.9)										
	7.2 (6.8)										
	7.1 (6.7)										
	7.0% (6.6)										
	6.9 (6.5)										
	6.8 (6.4)										
	6.7 (6.3)										
6.6 (6.2)											
6.5% (6.1)											
6.4 (6.0)											
6.3 (5.9)											
6.2 (5.8)											
6.1 (5.7)											
6.0% (5.6)											
5.9 (5.5)											
5.8 (5.4)											
5.7 (5.3)											
5.6% (5.2)											
5.5 (5.1)											
5.4 (5.0)											
5.3 ()											
5.2 (4.9)											
5.1 (4.8)											
5.0 (4.7)											
4.9 (4.6)											
糖代謝検査	HbA1c (%)	NGSP値									
		(JDS値)									
	血糖値 (mg/dl)	空腹時									
		食後 食後時間									

(科学的根拠に基づく糖尿病診療ガイドライン2013)

その他の検査など									